Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

and the points F_1, E, F_2, D are concyclic. The perpendicular bisectors of F_1F_2 and DE therefore intersect at C, the required center.

It may be observed that the second construction is a direct generalization of the one (when the asymptotes and vertex are given) found in all elementary text-books on Analytic Geometry. Would it not be well for the latter to contain the above simple constructions?

The University of Chicago, October, 1902.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ARITHMETIC.

161. Proposed by F. M. SHIELDS, Coopwood, Miss.

If 1 man, 1 boy, and 1 girl catch 1 trout, 1 perch, and one minnow in 5 minutes, and 1 man, 2 boys, and 3 girls catch 1 trout, 2 perch, and 3 minnows in 6 minutes, how many minutes will be required for 2 men, 3 boys, and 4 girls to catch 5 trout, 11 perch, and 17 minnows?

Remarks by B. F. FINKEL, A. M., M. Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

This problem is indeterminate. Let A_1, A_2, and A_3 be the activities of a man, boy, and girl, respectively; and E_1, E_2, and E_3 the work required to catch a trout, a perch, and a minnow, respectively. Then

$$5(A_1+A_2+A_3)=E_1+E_2+E_3 \ldots\ldots(1), \text{ and}$$
$$6(A_1+2A_2+3A_3)=E_1+2E_2+3E_3 \ldots\ldots(2).$$

From these two conditions it is required to find t so that the following condition is satisfied, viz:

$$t(2A_1+3A_2+4A_3)=5E_1+11E_2+17E_3.$$

This is clearly impossible in general. By assuming certain other conditions as, for example, the ratios of the activities, and the ratios of the work, a solution may be effected.

162. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

A trolly road is built between two towns, and it is found that the gross annual receipts amount to 20% of the original cost; the annual cost of repairs 2% of the original cost; and the working expense is $3000 in addition to 20% of the net receipts. After a year a second road is built at the same cost as the first and it is found that the gross re-